

Positive solutions of Robin problem for semilinear elliptic equations and a threshold result

Junhui Xie, Qiuyi Dai*, Huaxiang Hu
Department of Mathematics, Hunan Normal University
Changsha Hunan 410081, PR China

Abstract Let Ω be a bounded domain in R^n . The main topics of this paper are the following Robin problem for semilinear elliptic equation

$$\begin{cases} -\Delta u = u^p + f(x), & x \in \Omega, \\ u > 0, & x \in \Omega, \\ \frac{\partial u}{\partial \nu} + \beta u = 0, & x \in \partial\Omega, \end{cases} \quad (0.1)$$

and its parabolic version

$$\begin{cases} u_t - \Delta u = u^p + f(x), & (x, t) \in \Omega \times (0, T), \\ \frac{\partial u}{\partial \nu} + \beta u = 0, & (x, t) \in \partial\Omega \times [0, T), \\ u(x, 0) = u_0(x) \geq 0, & x \in \Omega. \end{cases} \quad (0.2)$$

We prove that for any given function $f(x)$ satisfies $(\mathcal{F})$ displayed in the introduction, there exists a positive number β_f^* such that problem (0.1) has no solution when $\beta \in (0, \beta_f^*)$, and has at least two solutions when $\beta > \beta_f^*$. Moreover, among all solutions of problem (0.1) there is a minimal one. Concerning problem (0.2), we prove that the minimal solution of problem (0.1) is stable, whereas, any other solution of problem (0.1) is an initial datum threshold for the existence and nonexistence of global solutions to problem (0.2). For more details, see Theorem 1.1-1.4 in the introduction.

Key words: Positive solution; Robin problem; threshold result; blow-up

1 Introduction

Let $n \geq 2$ and Ω be a bounded domain in R^n . Denoting by $\partial\Omega$ the boundary of Ω , we consider the following problem

$$\begin{cases} -\Delta u = u^p + f(x), & x \in \Omega, \\ u > 0, & x \in \Omega, \\ \frac{\partial u}{\partial \nu} + \beta u = 0, & x \in \partial\Omega, \end{cases} \quad (1.1)$$

where ν is the outward unit vector normal to $\partial\Omega$, $\beta \geq 0$ is a parameter and $f(x) \geq 0, f(x) \not\equiv 0$ is a given function in $C^1(\overline{\Omega})$.

Problem (1.1) has different names depending on the different values of the parameter β . It is called Dirichlet if $\beta = +\infty$, is called Neumann in the case $\beta = 0$ and is called Robin provided that $0 < \beta < +\infty$.

* Corresponding author. E-mail address: daiqiuyi@yahoo.com.cn (Q. Dai)
 This work is supported by NNSFC (Grant: No.10971061)

It is worth pointing out that problem (1.1) occurs in various branches of mathematical physics and biological models. Theoretically, the main topics in the study of problem (1.1) are to investigate the structure of solution sets and the stability or instability of its solutions.

In the case $\beta = 0$, it is trivial to see that problem (1.1) has no solution. In the case $\beta = +\infty$, there are many literature on problem (1.1). If $f(x) \equiv 0$, then problem (1.1) is reduced to the following problem

$$\begin{cases} -\Delta u = u^p, & x \in \Omega, \\ u > 0, & x \in \Omega, \\ u = 0, & x \in \partial\Omega. \end{cases} \quad (1.2)$$

It is well-known now that the existence of solutions to problem (1.2) strongly depends on the range of p . If $0 < p < 1$, then problem (1.2) has a unique solution, if $1 < p < \frac{n+2}{n-2}$, then problem (1.2) has at least one solution, if $p \geq \frac{n+2}{n-2}$ and Ω is a star shaped domain then problem (1.2) has no solution. Uniqueness and multiple results have also been obtained by many authors for the case $1 < p < \frac{n+2}{n-2}$ and some special domains (see for example [15, 3, 7]). A celebrate result given by Gidas and Spruck [11] says that the solution set of problem (1.2) is compact in $L^\infty(\Omega)$ when $1 < p < \frac{n+2}{n-2}$, though it may have many solutions. In 1992, Tarantello investigate the effect of inhomogeneous term $f(x)$ on the structure of solution set of problem (1.2), she considered in [22] the following problem

$$\begin{cases} -\Delta u = |u|^{p-1}u + f(x), & x \in \Omega, \\ u = 0, & x \in \partial\Omega, \end{cases} \quad (1.3)$$

and proved that problem (1.3) admits at least two solutions $u_0(x), u_1(x) \in H_0^1(\Omega)$ provided that $1 < p \leq \frac{n+2}{n-2}$, $f \not\equiv 0$ and

$$\int_{\Omega} f u dx \leq C_n \|\nabla u\|_{L^2(\Omega)}^{\frac{n+2}{2}} \quad \forall u \in H_0^1(\Omega),$$

where $C_n = \frac{4}{n-2}(\frac{n-2}{n+2})^{(n+2)/4}$. Furthermore $u_0(x) \geq 0, u_1(x) \geq 0$ when $f(x) \geq 0$. Roughly speaking, the appearance of the inhomogeneous term $f(x)$ increase the number of the solution. For more results on problem (1.3) see [4, 6, 9]

Compared with the Dirichlet Problem, there are few results on Robin problem. However, it is worth mentioning that Gu and Liu [18] studied the following problem

$$\begin{cases} -\Delta u = u^p, & x \in \Omega, \\ u > 0, & x \in \Omega, \\ \frac{\partial u}{\partial \nu} + \beta u = 0, & x \in \partial\Omega, \end{cases} \quad (1.4)$$

and proved that the solution set of problem (1.4) is compact in $L^\infty(\Omega)$. From the compactness of the solution set and the fixed point theorem, they also proved that problem (1.4) has at least one solution for any $0 < \beta < +\infty$ provided that $1 < p < \frac{n+2}{n-2}$. It is also worth pointing out that some existence, uniqueness and multiple results for problem (1.4) with $p \in (1, \frac{n+2}{n-2}]$ can be founded in [5, 10, 24].

In this paper, we focus our attention on the Robin problem with sub-critical Sobolev exponent p . Hence, we always assume that $0 < \beta < +\infty$ and $1 < p < \frac{n+2}{n-2}$ in the following paragraph. Our purpose are two-folds, one is to investigate the presence of the inhomogeneous term $f(x)$ how to change the structure of solution set of problem (1.4), the other is to investigate the stability and instability of solutions of problem (1.1). Concerning the first

issue, we find out that, unlike in the Dirichlet problem, the presence of the inhomogeneous term $f(x)$ in the Robin problem can increase, as well as, decrease the number of solutions subject to the range of the parameter β . Concerning the second issue, we find out that the minimal solution of (1.1) is stable and any other solution of problem (1.1) is a initial data threshold for the existence and nonexistence of global solutions to the parabolic version of problem (1.1).

To state our results rigorously, we introduce the following notations and conditions imposed on $f(x)$.

Let $h(x)$ be the unique solution of the following problem

$$\begin{cases} -\Delta h = 1, & x \in \Omega, \\ h = 0, & x \in \partial\Omega. \end{cases} \quad (1.5)$$

Setting

$$\begin{cases} M_h = \max_{x \in \Omega} h^p(x), \\ \Lambda = \left(\frac{1}{pM_h}\right)^{\frac{1}{p-1}}. \end{cases} \quad (1.6)$$

Then, a simple computation shows that

$$\Lambda - \Lambda^p M_h = \frac{p-1}{p} \left(\frac{1}{pM_h}\right)^{\frac{1}{p-1}} > 0. \quad (1.7)$$

For some results in this paper, the condition we impose on $f(x)$ can be read as

$$(\mathcal{F}) \quad f(x) \not\equiv 0 \quad \text{and} \quad 0 \leq f(x) < \frac{p-1}{p} \left(\frac{1}{pM_h}\right)^{\frac{1}{p-1}}.$$

The first result of this paper can be stated as

Theorem 1.1. For any given $f(x) \geq 0$ and $f(x) \not\equiv 0$. There exists a positive number β_f such that problem (1.1) has no solution for any $\beta \in (0, \beta_f)$.

The second result is about the existence of minimal solution and multiple result of problem (1.1). It can be presented as

Theorem 1.2. Assume that $f(x)$ satisfies $(\mathcal{F})$. Then there exists a positive number β_f^* such that

- (i) Problem (1.1) has no solution when $\beta \in (0, \beta_f^*)$.
- (ii) If $\beta \geq \beta_f^*$, then problem (1.1) has a minimal solution $U_\beta(x)$ in the sense that for any solution $u_\beta(x)$ of problem (1.1) we have $U_\beta(x) \leq u_\beta(x)$. Moreover U_β is decreasing with respect to the parameter β .
- (iii) If $\beta > \beta_f^*$, then problem (1.1) has at least two solutions.

Remark 1. Let $\Omega_R = \{x \in R^n \mid R < |x| < 1\}$, and $U_\beta(x)$ be the minimal solution of problem (1.1) with $\Omega = \Omega_R$. Setting $u(x) = v(x) + U_\beta(x)$, then $v(x)$ satisfies

$$\begin{cases} -\Delta v = (U_\beta(x) + v)^p - U_\beta^p(x), & x \in \Omega, \\ \frac{\partial v}{\partial \nu} + \beta v = 0, & x \in \partial\Omega. \end{cases} \quad (1.8)$$

As in [10], we can prove that for any fixed integer k , there exist at least k positive solutions to problem (1.8) provided that R is close enough to 1 and β is sufficiently large. As a result, problem (1.1) has at least $k+1$ solutions in this case.

Next, we consider the parabolic version of problem (1.1), that is, we consider

$$\begin{cases} u_t - \Delta u = u^p + f(x), & (x, t) \in \Omega \times (0, T), \\ \frac{\partial u}{\partial \nu} + \beta u = 0, & (x, t) \in \partial\Omega \times [0, T], \\ u(x, 0) = u_0(x) \geq 0, & x \in \Omega, \end{cases} \quad (1.9)$$

when $\beta = +\infty$ and $f(x) \equiv 0$, problem (1.9) is reduced to the following well studied problem

$$\begin{cases} u_t - \Delta u = u^p, & (x, t) \in \Omega \times (0, T), \\ u = 0, & (x, t) \in \partial\Omega \times [0, T], \\ u(x, 0) = u_0(x) \geq 0, & x \in \Omega. \end{cases} \quad (1.10)$$

The a priori bound and decaying estimate of global solutions of (1.10) can be found in [12] and [20, 21]. A threshold result of (1.10) under the condition that (1.2) has a unique solution was displayed in [2], similar result can also be found in [19]. Unfortunately, the uniqueness result is, in general, not true for problem (1.2) (see [15, 3, 17, 16]). It is worth pointing out that in the case $\Omega = \mathbb{R}^n$ many interesting results (including threshold result) of problem (1.10) can be found in [8, 24, 13, 14]. In this paper, we try to generalize the threshold result of problem (1.10) to problem (1.9). The main feature of our results is that we need not put uniqueness restriction on the steady state.

Theorem 1.3. For any given $f(x) \geq 0$ and $f(x) \not\equiv 0$, let β_f be the positive number obtained in Theorem 1.1. Then for any initial data $u_0(x) \geq 0$, the local solution $u(x, t, u_0)$ of problem (1.9) always blows up in finite time, that is, there exists $0 < T < +\infty$ such that

$$\lim_{t \rightarrow T^-} \max_{x \in \Omega} u(x, t, u_0) = +\infty.$$

Theorem 1.4. Assume that $f(x)$ satisfies $(\mathcal{F})$ and $1 < p < \frac{n+2}{n-2}$. Let β_f^* denote the positive number obtained in Theorem 1.2 and U_β be the minimal solution of problem (1.1). If $u_\beta(x)$ is an arbitrary solution of problem (1.1) which is distinct to U_β , then we have

- (i) For any initial data $u_0(x)$, problem (1.9) has no global solution when $0 < \beta < \beta_f^*$.
- (ii) If $\beta > \beta_f^*$, $0 \leq u_0(x) \leq u_\beta(x)$ and $u_0(x) \not\equiv u_\beta(x)$, then problem (1.9) has a global solution $u(x, t, u_0)$. Moreover,

$$\lim_{t \rightarrow +\infty} u(x, t, u_0) = U_\beta.$$

- (iii) If $\beta > \beta_f^*$, $u_0(x) \geq u_\beta(x)$ and $u_0(x) \not\equiv u_\beta(x)$, then the solution of problem (1.9) always blow up in finite time.

Remark 2. Roughly speaking. Theorem 1.4 implies that the minimal solution of problem (1.1) is stable, while any other solution of (1.1) is an initial datum threshold for the existence and nonexistence of global solution to problem (1.9).

Remark 3. In the case $f(x) \equiv 0$, problem (1.9) becomes

$$\begin{cases} u_t - \Delta u = u^p, & (x, t) \in \Omega \times (0, T), \\ \frac{\partial u}{\partial \nu} + \beta u = 0, & (x, t) \in \partial\Omega \times [0, T], \\ u(x, 0) = u_0(x) \geq 0, & x \in \Omega. \end{cases} \quad (1.11)$$

Concerning problem (1.11), we have

Theorem 1.5. Assume that $1 < p < \frac{n+2}{n-2}$. Let $U(x)$ be an arbitrary solution of problem (1.4). Then we have

(i) If $0 \leq u_0(x) \leq U(x)$ and $u_0(x) \not\equiv U(x)$, then problem (1.11) has a global solution $u(x, t; u_0)$, moreover,

$$\lim_{t \rightarrow +\infty} u(x, t; u_0) = 0.$$

(ii) If $u_0(x) \geq U(x)$ and $u_0(x) \not\equiv U(x)$, then the solution $u(x, t; u_0)$ of problem (1.11) always blow up in finite time.

Unlike in [2], our threshold result Theorem 1.5 need not put the uniqueness restriction on problem (1.4).

2 The proof of Theorem 1.1

This section is devoted to prove Theorem 1.1. To this end, we first prove some lemmas needed in the proof of Theorem 1.1 as follows.

Lemma 2.1. ([11]) If $u(x)$ is a solution of problem

$$\begin{cases} -\Delta u = u^p, & x \in R^n, \\ u \geq 0, & x \in R^n, \end{cases} \quad (2.1)$$

with $1 < p < \frac{n+2}{n-2}$, then $u(x) \equiv 0$.

Lemma 2.2. For $\beta \in (0, 1)$ small enough, there exists a positive constant C independent of β such that any solution $u = u_\beta$ of problem (1.1) satisfies $\|u_\beta\|_{C^{2,\alpha}(\bar{\Omega})} \leq C$.

Proof. By elliptic theory, it is enough to show that $\|u_\beta\|_{L^\infty(\bar{\Omega})} \leq C$. We argue by contradiction. Suppose that the conclusion is not true. Then there would exist a sequence $0 < \beta_j < \beta_f$ with $\beta_j \rightarrow 0$ as $j \rightarrow \infty$, a corresponding sequence of solutions $u_j = u_{\beta_j}$ of problem (1.1), and a sequence of points x_j in Ω such that

$$M_j = \|u_j\|_{L^\infty(\bar{\Omega})} = u_j(x_j) \rightarrow \infty \quad \text{as } j \rightarrow \infty. \quad (2.2)$$

Let us consider the auxiliary function

$$v_j(y) = M_j^{-1} u_j(x_j + M_j^{-\frac{p-1}{2}} y),$$

which is defined on $\Omega_j = M_j^{\frac{p-1}{2}} (\Omega - x_j)$.

It is easy to verify that $v_j(y)$ satisfies

$$\begin{cases} -\Delta v_j = v_j^p + M_j^{-p} f(x_j + M_j^{-\frac{p-1}{2}} y), & y \in \Omega_j, \\ v_j(0) = 1, \\ \frac{\partial v_j(y)}{\partial \nu} + \beta_j M_j^{-\frac{p-1}{2}} v_j(y) = 0, & y \in \partial\Omega_j, \\ 0 \leq v_j(y) \leq 1, & y \in \Omega_j. \end{cases} \quad (2.3)$$

We denote by D either the whole space R^n or the half space $R_+^n = \{x = (x_1, x_2, \dots, x_{n-1}, x_n) \in R^n \mid x_n > 0\}$. For any compact domain $K \subset D$, there exists a positive integer J such that $K \subset \Omega_j$ when $j > J$ due to $\Omega_j \rightarrow D$ as $j \rightarrow +\infty$. Noting that $0 \leq v_j(y) \leq 1$ and

$f(x) \in C^1(\overline{\Omega})$, it follows from the standard elliptic estimate that there exists a positive constant C independent of j such that $\|v_j\|_{C^{2,\alpha}(K)} \leq C$ with $\alpha \in (0, 1)$ for all $j > J$. By diagonal method, up to a subsequence, we may assume that v_j converges uniformly to a C^2 function v on any compact subset of D .

Since $M_j^{-p}f(x_j + M_j^{-\frac{p-1}{2}}y) \rightarrow 0$ as $j \rightarrow +\infty$, by sending j to $+\infty$ in problem (2.3), we see that v satisfies

$$\begin{cases} -\Delta v = v^p, & y \in D, \\ v \geq 0, \\ \frac{\partial v}{\partial \nu} = 0, & y \in \partial D \text{ if } D = R_+^n, \\ v(0) = 1. \end{cases} \quad (2.4)$$

If $D = R^n$, then $v(x)$ is a solution of

$$\begin{cases} -\Delta v = v^p, & y \in R^n, \\ v \geq 0, & y \in R^n, \\ v(0) = 1, \end{cases} \quad (2.5)$$

this contradicts to Lemma 2.1.

If $D = R_+^n$, we define

$$\tilde{v} = \tilde{v}(y_1, y_2, \dots, y_n) = \begin{cases} v(y_1, y_2, \dots, y_n), & y \in R_+^n; \\ v(y_1, y_2, \dots, y_{n-1}, -y_n), & y \in R_-^n; \end{cases}$$

then $\tilde{v}$ satisfies

$$\begin{cases} -\Delta \tilde{v} = \tilde{v}^p & y \in R^n, \\ \tilde{v} \geq 0, & y \in R^n, \\ \tilde{v}(0) = 1, \end{cases} \quad (2.6)$$

this also contradicts to Lemma 2.1. $\square$

Proof of Theorem 1.1. We argue by contradiction. Suppose that the conclusion of Theorem 1.1 is false. Then there exists a sequence $\beta_j \rightarrow 0^+$ as $j \rightarrow \infty$ such that problem (1.1) with $\beta = \beta_j$ has at least one positive solution u_{β_j} , by Lemma 2.2 and the standard elliptic estimate, there exists a positive constant C independent of j such that $\|u_{\beta_j}\|_{C^{2,\alpha}(\overline{\Omega})} \leq C$. Hence, up to a subsequence, we may assume that $u_{\beta_j} \rightarrow u$ in $C^2(\Omega)$ as $j \rightarrow \infty$ and u is a solution of the following problem

$$\begin{cases} -\Delta u = u^p + f(x), & x \in \Omega, \\ u \geq 0, & x \in \Omega, \\ \frac{\partial u}{\partial \nu} = 0, & x \in \partial\Omega. \end{cases} \quad (2.7)$$

However, it is easy to see that problem (2.7) has no solution since $f(x) \geq 0$ and $f(x) \not\equiv 0$. A contradiction. $\square$

3 The proof of Theorem 1.2

This section devotes to prove theorem 1.2. Since the proof is relatively long, we divide it into the following lemmas.

Lemma 3.1. Let $h(x)$ be the solution of problem (1.5), and $\varphi_\beta(x)$ be the solution of the problem

$$\begin{cases} -\Delta\varphi = 1, & x \in \Omega, \\ \frac{\partial\varphi}{\partial\nu} + \beta\varphi = 0, & x \in \partial\Omega. \end{cases} \quad (3.1)$$

Then $\varphi_\beta(x)$ is monotonically decreasing with respect to the parameter β and $\lim_{\beta \rightarrow +\infty} \varphi_\beta(x) = h(x)$.

Proof. Let $\beta_1 < \beta_2$, φ_{β_1} and φ_{β_2} be the solution of problem (3.1) with $\beta = \beta_1$ and β_2 respectively. If we set $w = \varphi_{\beta_1} - \varphi_{\beta_2}$, the w satisfies

$$\begin{cases} -\Delta w = 0, & x \in \Omega, \\ \frac{\partial w}{\partial\nu} + \beta_1 w \geq 0, & x \in \partial\Omega. \end{cases} \quad (3.2)$$

It follows from the maximum principle that $w \geq 0$ in Ω . Hence, $\varphi_\beta(x)$ is monotonically decreasing with respect to β . Noting that $\varphi_\beta(x) \geq 0$ for any $\beta > 0$ and $x \in \Omega$, we conclude that $\varphi_\beta(x)$ converges to some function. Passing to the limit in Problem (3.1) with $\beta \rightarrow +\infty$, we finally obtain

$$\lim_{\beta \rightarrow +\infty} \varphi_\beta(x) = h(x).$$

Lemma 3.2. If $f(x)$ satisfies $(\mathcal{F})$. Then there exists a positive number β_f^* such that

(i) Problem (1.1) has no solution when $\beta \in (0, \beta_f^*)$.

(ii) If $\beta \geq \beta_f^*$, then problem (1.1) has a minimal solution $U_\beta(x)$ in the sense that for any solution $u_\beta(x)$ of problem (1.1) we have $U_\beta(x) \leq u_\beta(x)$. Moreover U_β is strictly decreasing with respect to the parameter β .

Proof. Let φ_β be the solution of problem (3.1) and Λ be the constant given in (1.6). Setting $v_\beta = \Lambda\varphi_\beta$, it is easy to verify that $v_\beta(x)$ satisfies

$$\begin{cases} -\Delta v_\beta = -\Lambda\Delta\varphi_\beta = \Lambda, & x \in \Omega, \\ \frac{\partial v_\beta}{\partial\nu} + \beta v_\beta = 0, & x \in \partial\Omega. \end{cases} \quad (3.3)$$

By Lemma 3.1 and the assumption $(\mathcal{F})$, we have

$$\Lambda - \Lambda^p \max_{x \in \Omega} \varphi_\beta^p - f(x) \longrightarrow \Lambda - \Lambda^p M_h - f(x) > 0 \quad \text{when } \beta \longrightarrow +\infty.$$

Consequently, there exists a positive number β^* such that

$$\Lambda - \Lambda^p \varphi_\beta^p - f(x) \geq \Lambda - \Lambda^p \max_{x \in \Omega} \varphi_\beta^p - f(x) > 0,$$

for any $\beta > \beta^*$. This and (3.3) imply that $v_\beta = \Lambda\varphi_\beta$ is a super-solution of problem (1.1) when $\beta > \beta^*$. On the other hand, 0 is obviously a sub-solution of problem (1.1) and $0 < v_\beta(x)$. Hence, the sub and super-solution method implies that problem (1.1) has at least one solution $u_\beta(x)$ for any $\beta > \beta^*$. Moreover, $0 < u_\beta(x) \leq v_\beta(x)$.

Define

$$\beta_f^* = \inf\{\beta^* \in (0, +\infty) \text{ such that problem (1.1) has solution for any } \beta > \beta^*\}.$$

It is obvious that problem (1.1) has at least one solution for any $\beta > \beta_f^*$. Moreover, by a similar argument to that used in the proof of Lemma 2.2, we can conclude that for any $\beta \in (\beta_f^*, \beta_f^* + 1)$, there exist a positive constant C such that any solution $u_\beta(x)$ satisfies

$\|u_\beta\|_{C^{2,\alpha}(\overline{\Omega})} \leq C$. Hence, up to a subsequence, we may suppose that $u_\beta(x)$ converges to a function $u_{\beta_f^*}(x)$ in $C^2(\Omega)$ as $\beta \rightarrow \beta_f^* + 0$. Passing to the limit in problem (1.1) with $\beta \rightarrow \beta_f^* + 0$, we know that $u_{\beta_f^*}(x)$ is a solution of problem (1.1) with $\beta = \beta_f^*$. Hence, problem (1.1) has at least one solution for any $\beta \geq \beta_f^*$. Moreover, we know from theorem 1.1 that $\beta_f^* > 0$. Next, we prove that problem (1.1) has no solution for any $\beta \in (0, \beta_f^*)$ by contradiction. supposing that there exist a number $\beta_0 \in (0, \beta_f^*)$ such that problem (1.1) with $\beta = \beta_0$ has a solution u_{β_0} . Then for any $\beta > \beta_0$, u_{β_0} satisfies

$$\begin{cases} -\Delta u_{\beta_0}(x) = u_{\beta_0}^p(x) + f(x), & x \in \Omega, \\ \frac{\partial u_{\beta_0}(x)}{\partial \nu} + \beta u_{\beta_0}(x) > \frac{\partial u_{\beta_0}(x)}{\partial \nu} + \beta_0 u_{\beta_0}(x) = 0, & x \in \partial\Omega. \end{cases} \quad (3.4)$$

This implies that $u_{\beta_0}(x)$ is a super-solution of problem (1.1) for any $\beta > \beta_0$. On the other hand, 0 is a sub-solution of problem (1.1) and $0 < u_{\beta_0}(x)$. Hence, by the sub-and super-solution method, problem (1.1) has at least one solution for any $\beta > \beta_0$. This contradicts to the definition of β_f^* . Consequently, problem (1.1) has no solution for any $\beta \in (0, \beta_f^*)$.

To find a minimal solution of problem (1.1), we let $u_0(x) \equiv 0$ and $u_{j+1}(x)$ is the unique solution of the following problem

$$\begin{cases} -\Delta u_{j+1} = u_j^p + f(x), & x \in \Omega, \\ \frac{\partial u_{j+1}}{\partial \nu} + \beta u_{j+1} = 0, & x \in \partial\Omega. \end{cases} \quad (3.5)$$

Then it is easy to see that $\{u_j(x)\}$ is a monotonically increasing sequence. Moreover, if $u_\beta(x)$ is a solution of problem (1.1), we also have $u_j(x) \leq u_\beta(x)$ for any $x \in \Omega$ and any $j > 0$. Hence, $\{u_j(x)\}$ is convergence when j tends to the infinity. Let

$$U_\beta(x) = \lim_{j \rightarrow +\infty} u_j(x).$$

Then, $U_\beta(x)$ is obviously the minimal solution of problem (1.1).

To prove that $U_\beta(x)$ is monotonically decreasing with respect to the parameter β , we assume that $\beta_2 > \beta_1$ and $U_{\beta_2}(x)$ and $U_{\beta_1}(x)$ are the minimal solution of problem (1.1) with $\beta = \beta_2$ and $\beta = \beta_1$ respectively. Then, a simple computation shows that $U_{\beta_1}(x)$ satisfies

$$\begin{cases} -\Delta U_{\beta_1}(x) = U_{\beta_1}^p(x) + f(x), & x \in \Omega, \\ \frac{\partial U_{\beta_1}(x)}{\partial \nu} + \beta_2 U_{\beta_1}(x) > \frac{\partial U_{\beta_1}(x)}{\partial \nu} + \beta_1 U_{\beta_1}(x) = 0, & x \in \partial\Omega. \end{cases} \quad (3.6)$$

Thus, $U_{\beta_1}(x)$ is a super-solution of problem (1.1) with $\beta = \beta_2$. By the definition of $U_{\beta_2}(x)$, we have $U_{\beta_2}(x) \leq U_{\beta_1}(x)$ for any $x \in \Omega$. Moreover, by strong maximum principle, we have $U_{\beta_2}(x) < U_{\beta_1}(x)$ for any $x \in \Omega$. That is, $U_\beta(x)$ is strictly decreasing with respect to the parameter β . $\square$

For any $\beta > \beta_f^*$, by making use of the minimal solution of problem (1.1), we can decompose any solution $u_\beta(x)$ of problem (1.1) as

$$u_\beta(x) = v(x) + U_\beta(x),$$

where $v(x)$ satisfies

$$\begin{cases} -\Delta v = (U_\beta(x) + v)^p - U_\beta^p(x), & x \in \Omega, \\ \frac{\partial v}{\partial \nu} + \beta v = 0, & x \in \partial\Omega. \end{cases} \quad (3.7)$$

To find another solution of problem (1.1), we have only to find a positive solution of problem (3.7). To do this, we represent $(U_\beta(x) + v)^p - U_\beta^p(x)$ as

$$(U_\beta(x) + v)^p - U_\beta^p(x) = pU_\beta^{p-1}(x) + g(x, v).$$

Then, $g(x, v)$ satisfies

$$(g_1) \lim_{v \rightarrow 0} \frac{g(x, v)}{v} = 0 \text{ uniformly on } \Omega;$$

$$(g_2) \lim_{v \rightarrow 0} \frac{g(x, v)}{v^p} = 0 \text{ uniformly on } \Omega \text{ with } 1 < p < \frac{n+2}{n-2};$$

$$(g_3) g(x, 0) = 0 \text{ and } g(x, v) > 0 \text{ for } v > 0;$$

(g_4) there exists some $\theta > 2$ and $M > 0$ such that $0 < \theta G(x, v) \leq vg(x, v)$ for all $v \geq M$ and $x \in \Omega$.

The following lemma is crucial for finding positive solutions of problem (3.7).

Lemma 3.3. The first eigenvalue $\lambda_1(\Omega)$ of the following eigenvalue problem

$$\begin{cases} -\Delta\phi - pU_\beta^{p-1}\phi = \lambda\phi, & x \in \Omega, \\ \frac{\partial\phi}{\partial\nu} + \beta\phi = 0, & x \in \partial\Omega, \end{cases} \quad (3.8)$$

is positive.

Proof. Let $\phi_1(x)$ be the first eigenfunction of problem (3.8). It is well known that $\phi_1(x)$ can be chosen so that $\phi_1(x) > 0$ for any $x \in \Omega$. Let $\beta_1 < \beta$, $U_{\beta_1}(x)$ and $U_\beta(x)$ be the minimal solution of problem (1.1) with parameter β_1 and β in boundary condition respectively. Then, it follows from lemma 3.2 that

$$U_{\beta_1}(x) > U_\beta(x) \text{ for } x \in \Omega.$$

Setting $v(x) = U_{\beta_1}(x) - U_\beta(x)$, then $v(x) > 0$ for any $x \in \Omega$. Furthermore, a simple calculation implies that $v(x)$ satisfies

$$\begin{cases} -\Delta v = U_{\beta_1}^p - U_\beta^p > pU_\beta^{p-1}v, & x \in \Omega, \\ \frac{\partial v}{\partial\nu} + \beta v = 0, & x \in \partial\Omega. \end{cases} \quad (3.9)$$

From this, we have

$$\beta \int_{\partial\Omega} v\phi_1 ds + \int_{\Omega} \nabla v \bullet \nabla \phi_1 dx - p \int_{\Omega} U_\beta^{p-1} v\phi_1 dx > 0. \quad (3.10)$$

On the other hand, by the equations satisfied by $\phi_1(x)$, we can deduce that

$$\beta \int_{\partial\Omega} v\phi_1 ds + \int_{\Omega} \nabla v \bullet \nabla \phi_1 dx - p \int_{\Omega} U_\beta^{p-1} v\phi_1 dx = \lambda_1(\Omega) \int_{\Omega} v\phi_1 dx. \quad (3.11)$$

Combining (3.10) with (3.11), we obtain

$$\lambda_1(\Omega) \int_{\Omega} v\phi_1 dx > 0.$$

Since $\int_{\Omega} v\phi_1 dx > 0$, we reach $\lambda_1(\Omega) > 0$. $\square$

Proof of Theorem 1.2. The proof of theorem 1.2 (i) is similar to that of theorem 1.1. The conclusion (ii) of theorem 1.2 comes from lemma 3.2. To prove (iii) of theorem 1.2, let $H^1(\Omega)$ be the usual Sobolev space endowed with the usual norm

$$\|u\| = \sqrt{\|u\|_{L^2(\Omega)}^2 + \|\nabla u\|_{L^2(\Omega)}^2}$$

From lemma 3.3, we can easily see that for $\beta > \beta_f^*$

$$\|u\|_* = \sqrt{\beta \int_{\partial\Omega} v^2 ds + \int_{\Omega} |\nabla v|^2 dx - p \int_{\Omega} U_{\beta}^{p-1} v^2 dx}$$

is a norm equivalent to $\|u\|$. Denote by $H_*^1(\Omega)$ the function space $H^1(\Omega)$ endowed with the norm $\|\bullet\|_*$. Then $H_*^1(\Omega)$ is obviously a Banach space. Consider the functional

$$\begin{aligned} I(v) &= \frac{1}{2} \int_{\Omega} |\nabla v|^2 dx - \frac{1}{2} p \int_{\Omega} U_{\beta}^{p-1} v^2 dx + \frac{\beta}{2} \int_{\partial\Omega} v^2 ds - \int_{\Omega} G(x, v^+) dx \\ &= \frac{1}{2} \|u\|_*^2 - \int_{\Omega} G(x, v^+) dx \end{aligned}$$

which is defined on $H_*^1(\Omega)$, where $v^+(x) = \max\{0; v(x)\}$. By making use of $(g_1) \sim (g_4)$ satisfied by $g(x, v)$, we can verify that $I(v)$ satisfies all desired conditions in the Mountain-Pass Theorem with PS-condition (see[9] for more details). Thus, Mountain-Pass Theorem with PS-condition [1] assures that problem (3.7) has at least one positive solution for $\beta > \beta_f^*$. Consequently, problem (1.1) has at least two solution for $\beta > \beta_f^*$. This completes the proof of theorem 1.2. $\square$

4 A priori bound for the global solution of problem (1.9)

In this section, we derive an a priori bound for global solution of problem (1.9). More precisely, we will prove the following theorem.

Theorem 4.1. Assume that $1 < p < \frac{n+2}{n-2}$, and $u(x, t)$ is a classical solution of problem (1.9) in $\Omega \times (0, +\infty)$ with $0 \leq u_0(x) \in C(\Omega)$. Then there exists a positive constant M depending only on $\max_{x \in \Omega} u_0(x)$ and $\sup_{x \in \Omega} f(x)$ such that $u(x, t) \leq M$ for any $(x, t) \in \Omega \times (0, +\infty)$.

When $\beta = +\infty$ and $f(x) \equiv 0$, theorem 4.1 was proved by Giga in [12]. Apart from some minor modification, the proof of theorem 4.1 can be adapted from [12] line by line. However, for the reader's convenience and the need of the sequel paragraph we give a sketch proof here.

Lemma 4.1. Let $u(x, t)$ be a nontrivial nonnegative solution of problem (1.9) in $Q = \Omega \times [0, T)$. Suppose that there exists a positive number N independent of T such that $u(x, t)$ satisfies

$$\int_0^T \int_{\Omega} |u_t|^2 < N, \quad (4.1)$$

and that for a given $t_0 > 0$, $\sup_{x \in \Omega \times [0, T)} u(x, t)$ is attained in $\Omega \times (t_0, T)$. Then there is a constant A depending only on N and t_0 such that $u(x, t) \leq A$ for any $(x, t) \in Q$.

Proof. Suppose the lemma is false. Then there exists a sequence of solutions $u_k(x, t)$ of problem (1.9) with $T = T_k > 0$ and a sequence of points $(x_k, t_k) \in \Omega \times (t_0, T_k)$ such that

$$M_k = \sup_{(x, t) \in \Omega \times [0, T_k)} u_k(x, t) = u_k(x_k, t_k) \rightarrow \infty, \text{ as } k \rightarrow \infty. \quad (4.2)$$

and

$$\int_0^{T_k} \int_{\Omega} |u_t|^2 < N. \quad (4.3)$$

Let λ_k be a sequence of positive numbers such that

$$\lambda_k^{\frac{2}{p-1}} M_k = 1.$$

Then $\lambda_k \rightarrow 0$ as $k \rightarrow \infty$ due to $M_k \rightarrow +\infty$.

Define a sequence of function $v_k(y, s)$ by

$$v_k(y, s) = \lambda_k^{\frac{2}{p-1}} u_k(x_k + \lambda_k y, t_k + \lambda_k^2 s). \quad (4.4)$$

Let $Q_k = \Omega_k \times (-\lambda_k^{-2} t_k, 0]$ with $\Omega_k = \lambda_k^{-1}(\Omega - x_k) = \{y \mid y = \lambda_k^{-1}(x - x_k), x \in \Omega\}$, and $\partial Q_k = \partial \Omega_k \times [-\lambda_k^{-2} t_k, 0]$. Then, it is easy to verify that $v_k(y, s)$ satisfies

$$\begin{cases} v_{ks} - \Delta_y v_k = v_k^p + \lambda_k^{\frac{2p}{p-1}} f(x_k + \lambda_k y), & (y, s) \in Q_k, \\ \frac{\partial v_k}{\partial \nu} + \lambda_k \beta v_k = 0, & (y, s) \in \partial Q_k, \\ v_k(y, s) \leq 1, & (y, s) \in Q_k, \\ v_k(0, 0) = 1. \end{cases} \quad (4.5)$$

Let $d_k = \text{dist}(x_k, \partial \Omega)$ denote the distance from x_k to $\partial \Omega$, and $R_a^n = \{y = (y_1, y_2, \dots, y_n) \in R^n \mid y_n > -a\}$. Since $t_k > t_0$, it is easy to see that up to a subsequence, there hold

$$Q_k \rightarrow R^n \times (-\infty, 0] \quad \text{or} \quad Q_k \rightarrow R_a^n \times (-\infty, 0]$$

subject to $\lambda_k^{-1} d_k \rightarrow +\infty$ or $\lambda_k^{-1} d_k \rightarrow a$ respectively. Hence, our proof are divided into the following two cases.

Case (i). If $\lambda_k^{-1} d_k \rightarrow +\infty$, then $Q_k \rightarrow R^n \times (-\infty, 0]$. Consequently, for any parabolic cylinder $Q(r) = \{(y, s) \in R^{n+1} \mid |y| < r, s \in (-r^2, 0]\}$ with radius r in R^{n+1} , there exists a positive integer K such that $Q(r) \subset Q_k$ for all $k > K$. Since $v_k(y, s) \leq 1$ in Q_k , it follows from the parabolic L^q theory [25] that $v_k(y, s)$ is uniformly bounded in $W_q^{2,1}(Q(r))$ for any $q > n$. Hence, by the diagonal method, we can choose a subsequence of $v_k(y, s)$, still denote it by $v_k(y, s)$, and a function $v \geq 0$ defined on $R^n \times (-\infty, 0]$ such that $v_k(y, s)$ converges uniformly to v in any parabolic cylinder $Q(r)$. Furthermore, by taking another subsequence if necessary, we may assume that $v_{ks}(y, s)$ converges to v_s in $L^2(Q(r))$. Passing to the limit in problem (4.5) with $k \rightarrow +\infty$, and noting that $\lambda_k^{\frac{2p}{p-1}} f(x_k + \lambda_k y) \rightarrow 0$ as $k \rightarrow +\infty$, we see that v solves

$$\begin{cases} v_s - \Delta v = v^p, & (y, s) \in R^n \times (-\infty, 0], \\ v(y, s) \geq 0, & (y, s) \in R^n \times (-\infty, 0], \\ v(0, 0) = 1. \end{cases} \quad (4.6)$$

Thanks to (4.3), we have $v_s \equiv 0$ in $R^n \times (-\infty, 0)$. In fact, a simple computation shows that

$$\begin{aligned} \int_{Q(r)} |v_{ks}|^2 dy ds &= \lambda_k^{\frac{4}{p-1} - n + 2} \int_{t_k - \lambda_k^2 r^2}^{t_k} \int_{|x - x_k| < \lambda_k r} |u_t|^2 dx dt \\ &\leq \lambda_k^{\frac{4}{p-1} - n + 2} \int_0^{T_k} \int_{\Omega} |u_t|^2 dx dt \\ &\leq \lambda_k^{\frac{4}{p-1} - n + 2} N. \end{aligned} \quad (4.7)$$

Since $\frac{4}{p-1} - n + 2 > 0$ due to $p \in (1, \frac{n+2}{n-2})$, we have

$$\int_{Q(r)} |v_{ks}|^2 dy ds \rightarrow 0.$$

This yields $v_s = 0$ in $Q(r)$ because v_{ks} converges weakly to v_s in $L^2(Q(r))$ and the norm is lower semi-continuous under weak convergence. Noting further that r is arbitrary, v_s vanishes identically in $R^n \times (-\infty, 0)$. Hence, v is independent of s , and $v \geq 0$ satisfies

$$\begin{cases} -\Delta v = v^p, & \text{in } R^n, \\ v(0) = 1. \end{cases} \quad (4.8)$$

This contradicts Lemma 2.1.

Case (ii). If $\lambda_k^{-1} d_k \rightarrow a$, then $Q_k \rightarrow R_a^n \times (-\infty, 0]$. Let $Q_a = R_a^n \times (-\infty, 0]$. Then, for any parabolic cylinder $Q(r)$ in R^{n+1} , there exists a positive integer K such that $Q(r) \cap Q_a \subset Q_k$ for all $k > K$. By making use of parabolic L^q regularity theory up to the boundary, as in the case (i) we can choose a subsequence of v_k , still denote it by v_k , such that v_k converges uniformly in $Q(r) \cap Q_a$ to a function v defined on Q_a and such that v satisfies

$$\begin{cases} -\Delta v = v^p, & y \in \{y_n > -a\}, \\ \frac{\partial v}{\partial \nu} = 0, & y \in \{y_n = -a\}, \\ v(0) = 1. \end{cases} \quad (4.9)$$

Changing coordinates by $y^* = (y_1, y_2, \dots, y_n + a)$, then $\hat{v}(y^*) = v(y)$ solves

$$\begin{cases} -\Delta \hat{v} = \hat{v}^p, & y^* \in R_+^n, \\ \frac{\partial \hat{v}}{\partial \nu} = 0, & y^* \in \partial R_+^n, \\ \hat{v}(0, a) = 1. \end{cases} \quad (4.10)$$

Define

$$\tilde{v}(y^*) = \begin{cases} \hat{v}(y_1^*, y_2^*, \dots, y_{n-1}^*, y_n^*), & y^* \in R_+^n; \\ \hat{v}(y_1^*, y_2^*, \dots, y_{n-1}^*, -y_n^*), & y^* \in R_-^n; \end{cases} \quad (4.11)$$

Then $\tilde{v}$ satisfies

$$\begin{cases} -\Delta \tilde{v} = \tilde{v}^p, & y^* \in R^n, \\ \tilde{v}(0, a) = 1. \end{cases} \quad (4.12)$$

This also contradicts Lemma 2.1. $\square$

Proof of Theorem 4.1. Assume that $u_0 \geq 0$ and $u_0(x) \not\equiv 0$. We first note that there are constant B , $t' > 0$ depending only on $\sup_{x \in \Omega} u_0(x)$, $|\Omega|$ and $\sup_{x \in \Omega} f(x)$ such that (see [25])

$$\sup_{0 \leq \tau \leq 2t'} \sup_{\Omega} u(x, t) \leq B, \quad \int_{\Omega} |\nabla u|^2(x, t') dx \leq B. \quad (4.13)$$

Let

$$E(t) = E[u] = \frac{1}{2} \int_{\Omega} |\nabla u|^2 dx + \frac{\beta}{2} \int_{\partial \Omega} u^2 ds - \frac{1}{p+1} \int_{\Omega} u^{p+1} dx - \int_{\Omega} f u dx \quad (4.14)$$

be the energy associated to problem (1.9). Then, it is easy to check that the following energy identities hold

$$\frac{d}{dt} \int_{\Omega} |u|^2 dx = -4E(t) + \frac{2(p-1)}{p+1} \int_{\Omega} u^{p+1} - 2 \int_{\Omega} f u dx, \quad (4.15)$$

$$\int_{\Omega} |u_t|^2 dx = -\frac{d}{dt} E(t). \quad (4.16)$$

By Holder and Young's inequality, we have

$$\begin{aligned} 2 \int_{\Omega} f u dx &\leq 2 \left(\int_{\Omega} u^{p+1} dx \right)^{\frac{1}{p+1}} \left(\int_{\Omega} f^{\frac{p+1}{p}} dx \right)^{\frac{p}{p+1}} \\ &\leq \frac{p-1}{p+1} \int_{\Omega} u^{p+1} dx + \left(\frac{2^{p+1}p}{p-1} \right)^{\frac{1}{p}} \int_{\Omega} f^{\frac{p+1}{p}} dx. \end{aligned} \quad (4.17)$$

It follows from (4.15) and (4.17) that

$$\frac{d}{dt} \int_{\Omega} |u|^2 dx \geq -4E(t) - \left(\frac{2^{p+1}p}{p-1} \right)^{\frac{1}{p}} \int_{\Omega} f^{\frac{p+1}{p}} dx + \left(\frac{p-1}{p+1} \right) \int_{\Omega} u^{p+1} dx. \quad (4.18)$$

Identity (4.16) says that the energy $E(t)$ should decrease. This together with (4.18) shows that for any $t > 0$, we have

$$E(t) \geq -\frac{1}{4} \left(\frac{2^{p+1}p}{p-1} \right)^{\frac{1}{p}} \int_{\Omega} f^{\frac{p+1}{p}} dx, \quad (4.19)$$

since otherwise the solution must blow up in finite time.

Integrating (4.16) over (t', T) gives

$$\begin{aligned} \int_{t'}^T \int_{\Omega} |u_t|^2 dx dt &= E(t') - E(T) \\ &\leq E(t') + \frac{1}{4} \left(\frac{2^{p+1}p}{p-1} \right)^{\frac{1}{p}} \int_{\Omega} f^{\frac{p+1}{p}} dx \\ &\leq \frac{B}{2} + \frac{1}{4} \left(\frac{2^{p+1}p}{p-1} \right)^{\frac{1}{p}} \int_{\Omega} f^{\frac{p+1}{p}} dx. \end{aligned} \quad (4.20)$$

Applying Lemma 4.1 with $Q = \Omega \times [t', T)$, we know that any solution in $\Omega \times [t', T)$ which attains its maximum outside $\Omega \times [t', 2t')$ is bounded from above by a constant M which is independent of T . By (4.13) we know that a solution which take its maximum inside $\Omega \times [t', 2t')$ is dominated by B . Hence in any case

$$u(x, t) \leq M + B \quad \text{for any } (x, t) \in \Omega \times [t', \infty). \quad (4.21)$$

Combining (4.13) with (4.21), we reach the desired conclusion of Theorem 4.1. $\square$

5 The proof of Theorem 1.3

This section devotes to prove theorem 1.3. So, we always assume that $\beta < \beta_f$ with β_f being the number determined in theorem 1.1. If theorem 1.3 is false, then there is a initial data $u_0(x)$ such that problem (1.9) with initial data $u_0(x)$ has a global solution $u(x, t; u_0)$. For simplicity, we denote $u(x, t; u_0)$ by $u(t)$ if no confusion arise. Let $E(t)$ be the energy associated to $u(t)$, that is

$$E(t) = \frac{1}{2} \int_{\Omega} |\nabla u(t)|^2 dx + \frac{\beta}{2} \int_{\partial\Omega} u^2(t) ds - \frac{1}{p+1} \int_{\Omega} u^{p+1}(t) dx - \int_{\Omega} f(x) u(t) dx. \quad (5.1)$$

A similar argument to that used in the proof of theorem 4.1 shows that for any $t > 0$,

$$E(t) \geq -\frac{1}{4} \left(\frac{2^{p+1}p}{p-1} \right)^{\frac{1}{p}} \int_{\Omega} f^{\frac{p+1}{p}} dx \quad (5.2)$$

and

$$\int_{\Omega} |u_t(t)|^2 dx = -\frac{d}{dt} E(t). \quad (5.3)$$

This yields

$$\int_0^\infty \int_\Omega |u_t(t)|^2 dx dt \leq E(u_0) + \frac{1}{4} \left(\frac{2^{p+1}p}{p-1} \right)^{\frac{1}{p}} \int_\Omega f^{\frac{p+1}{p}} dx < +\infty. \quad (5.4)$$

Thus, we can pick up a sequence $\{t_j\}$ such that $t_j \rightarrow \infty$ and $\|u_t(t_j)\|_{L^2(\Omega)} \rightarrow 0$ as $j \rightarrow +\infty$. Moreover, by theorem 4.1, we have $u(t_j) \leq M$ for some positive constant M independent of j .

Multiplying the first equation in problem (1.9) by $u(t_j)$ and integrating over Ω yields:

$$\begin{aligned} \beta \int_{\partial\Omega} u^2(t_j) dx + \int_\Omega |\nabla u(t_j)|^2 dx &= \int_\Omega u^{p+1}(t_j) dx + \int_\Omega f(x) u(t_j) dx - \int_\Omega u(t_j) u_t(t_j) dx \\ &\leq \int_\Omega u^{p+1}(t_j) dx + \int_\Omega f(x) u(t_j) dx + \int_\Omega |u(t_j) u_t(t_j)| dx \\ &\leq M^{p+1} |\Omega| + M \|f\|_{L^1(\Omega)} + M |\Omega|^{\frac{1}{2}} \|u(t_j)\|_{L^2(\Omega)}. \end{aligned} \quad (5.5)$$

Taking $\|u_t(t_j)\|_{L^2(\Omega)} \rightarrow 0$ into account, we arrive

$$\|u_j\|_{H^1(\Omega)} \leq M^{p+1} |\Omega| + M \|f\|_{L^1(\Omega)} + O(1).$$

From this and the weak compactness of $H^1(\Omega)$, we may assume, up to a subsequence, that for some function $u(x) \in H^1(\Omega)$

$$\begin{aligned} u(t_j) &\rightharpoonup u(x) \quad \text{weakly in } H^1(\Omega) \\ u(t_j) &\rightarrow u(x) \quad \text{strongly in } L^{p+1}(\Omega). \end{aligned} \quad (5.6)$$

For any given $\varphi(x) \in H^1(\Omega)$, multiplying the first equation in problem (1.9) by $\varphi(x)$ and integrating on Ω yields

$$\begin{aligned} \int_\Omega \varphi u_t(t_j) dx &= \int_\Omega \varphi \Delta u(t_j) dx + \int_\Omega u^p(t_j) \varphi dx + \int_\Omega f \varphi dx \\ &= -\beta \int_{\partial\Omega} u(t_j) \varphi ds - \int_\Omega \nabla u(t_j) \bullet \nabla \varphi dx + \int_\Omega u^p(t_j) \varphi dx + \int_\Omega f \varphi dx. \end{aligned} \quad (5.7)$$

Thus

$$\int_\Omega \varphi u_t(t_j) dx = -\beta \int_{\partial\Omega} u(t_j) \varphi ds - \int_\Omega \nabla u(t_j) \bullet \nabla \varphi dx + \int_\Omega u^p(t_j) \varphi dx + \int_\Omega f \varphi dx. \quad (5.8)$$

Taking (5.6) and $\|u_t(t_j)\|_{L^2(\Omega)} \rightarrow 0$ into account, by passing to the limit in (5.8) we arrive

$$\beta \int_{\partial\Omega} u(x) \varphi ds + \int_\Omega \nabla u(x) \bullet \nabla \varphi dx = \int_\Omega u^p(x) \varphi dx + \int_\Omega f \varphi dx. \quad (5.9)$$

This implies that $u(x)$ is a weak solution of problem (1.1). By standard regularity theory of elliptic equations, $u(x)$ is also a classical solution of problem (1.1). This contradicts theorem 1.1 and the proof of theorem 1.3 is completed. $\square$

6 The Proof of Theorem 1.4

We give a proof of theorem 1.4 in this section. To this end, we need the following lemmas first.

Lemma 6.1. Let b is a positive constant, $p > 1$ and $f(t) = ((t+b)^p - b^p)/t$. Then $f(t)$ is monotonically increasing in $(0, +\infty)$.

Proof. An easy computation yields

$$f'(t) = \frac{pt(t+b)^{p-1} - (t+b)^p + b^p}{t^2}.$$

By mean value theorem, there exists $\xi \in (0, t)$ such that

$$(t + b)^p - b^p = pt(\xi + b)^{p-1}.$$

hence

$$f'(t) = \frac{pt[(t + b)^{p-1} - (\xi + b)^{p-1}]}{t^2}.$$

Noting $p > 1$, we have $f'(t) \geq 0$ for any $t > 0$. $\square$

Lemma 6.2. Assume that $p > 1$ and $a > b > 0$. If $f(a) = a^p - pab^{p-1} + pb^p - b^p$, $g(a) = a^p - pa^p + pa^{p-1}b - b^p$, then $f(a) > 0$, $g(a) < 0$.

Proof. Since $f'(a) = p(a^{p-1} - b^{p-1}) > 0$ and $f(b) = 0$, we have $f(a) > 0$. Similarly, we can prove $g(a) < 0$. $\square$

Lemma 6.3. Assume that $p > 1$ and $a > b > 0$. If $F(\eta) = \eta(a^p - b^p) + b^p - [\eta(a - b) + b]^p$, then $F(\eta) > 0$ for $0 < \eta < 1$, and $F(\eta) < 0$ for $\eta > 1$.

Proof. An easy computation shows

$$\begin{cases} F'(\eta) = a^p - b^p - p(a - b)[\eta(a - b) + b]^{p-1}, \\ F''(\eta) = -p(p - 1)(a - b)^2[\eta(a - b) + b]^{p-2} < 0. \end{cases} \quad (6.1)$$

Hence, $F'(\eta)$ is strictly decreasing in $[0, +\infty)$. By lemma 6.2, we have

$$\begin{cases} F'(0) = a^p - pab^{p-1} + pb^p - b^p = f(a) > 0, \\ F'(1) = a^p - pa^p + pa^{p-1}b - b^p = g(a) < 0. \end{cases} \quad (6.2)$$

This implies that there exists a unique point $\xi \in (0, 1)$ such that $F'(\xi) = 0$. Hence

$$\begin{cases} F'(\eta) > 0 & \text{for } 0 < \eta \leq \xi, \\ F'(\eta) < 0 & \text{for } \xi < \eta \leq 1. \end{cases} \quad (6.3)$$

Consequently

$$\begin{cases} F(\eta) > F(0) = 0 & \text{for } 0 < \eta \leq \xi, \\ F(\eta) > F(1) = 0 & \text{for } \xi < \eta < 1, \\ F(\eta) < F(1) = 0 & \text{for } \eta > 1. \end{cases} \quad (6.4)$$

This is the desired conclusion of lemma 6.3. $\square$

The following lemma is important for our proof of theorem 1.4.

Lemma 6.4. If $u_1(x)$ and $u_2(x)$ are any two distinct solutions of problem (1.1) which are also different from the minimal one, then $u_1(x)$ and $u_2(x)$ must intersect somewhere.

Proof. Since $u_1(x)$ and $u_2(x)$ are solutions of problem (1.1) which are different from the minimal solution $U_\beta(x)$, we can decompose $u_1(x)$ and $u_2(x)$ as the following

$$u_1(x) = v_1(x) + U_\beta(x).$$

$$u_2(x) = v_2(x) + U_\beta(x).$$

Moreover, it follows from strong maximum principle that $v_1(x)$ and $v_2(x)$ are positive in $\overline{\Omega}$. By making use of equations satisfied by $u_1(x)$ and $u_2(x)$, it is easy to check that $v_1(x)$ and $v_2(x)$ solve the following problem respectively

$$\begin{cases} -\Delta v_1 = (U_\beta(x) + v_1)^p - U_\beta^p(x) & x \in \Omega \\ \frac{\partial v_1}{\partial \nu} + \beta v_1 = 0 & x \in \partial\Omega \end{cases} \quad (6.5)$$

$$\begin{cases} -\Delta v_2 = (U_\beta(x) + v_2)^p - U_\beta^p(x) & x \in \Omega \\ \frac{\partial v_2}{\partial \nu} + \beta v_2 = 0 & x \in \partial\Omega \end{cases} \quad (6.6)$$

From these, we can derive

$$\int_{\Omega} v_2[(U_\beta(x) + v_1)^p - U_\beta^p(x)]dx = \int_{\Omega} v_1[(U_\beta(x) + v_2)^p - U_\beta^p(x)]dx. \quad (6.7)$$

That is

$$\int_{\Omega} v_1 v_2 \left[\frac{(U_\beta(x) + v_1)^p - U_\beta^p(x)}{v_1} - \frac{(U_\beta(x) + v_2)^p - U_\beta^p(x)}{v_2} \right] dx = 0. \quad (6.8)$$

From this and the conclusion of lemma 6.1, we can easily see the validity of lemma 6.4. $\square$

Proof of Theorem 1.4. The proof of theorem 1.4 (i) is the same as that of theorem 1.3. To prove theorem 1.4 (ii) and (iii), we let $u_\beta(x)$ be a solution of problem (1.1) which is distinct to the minimal solution $U_\beta(x)$. Then, by the strong maximum principle, we have

$$U_\beta(x) < u_\beta(x) \quad \text{for any } x \in \Omega. \quad (6.9)$$

If the initial datum $0 \leq u_0(x) \leq u_\beta(x)$ and $u_0(x) \not\equiv u_\beta(x)$, then problem (1.9) has a global solution $u(x, t; u_0)$. Moreover, it follows from the strong comparison principle that $u(x, t; u_0)$ satisfies

$$u(x, t; u_0) < u_\beta(x) \quad \text{for any } (x, t) \in \bar{\Omega} \times (0, +\infty). \quad (6.10)$$

To prove $\lim_{t \rightarrow +\infty} u(x, t; u_0) = U_\beta(x)$, we may assume, by replacing $u_0(x)$ with $u(x, T; u_0)$ for some $T > 0$ if necessary, that $u_0(x) \leq \eta u_\beta(x)$ for some $0 < \eta < 1$. Let $h_\beta(x) = \eta(u_\beta(x) - U_\beta(x)) + U_\beta(x)$. Then, by making use of lemma 6.3 with $0 < \eta < 1$, we can verify that $h_\beta(x)$ satisfies

$$\begin{cases} -\Delta h_\beta > h_\beta^p + f(x), & x \in \Omega, \\ h_\beta > 0, & x \in \Omega, \\ \frac{\partial h_\beta}{\partial \nu} + \beta h_\beta = 0, & x \in \partial\Omega. \end{cases} \quad (6.11)$$

This implies that $h_\beta(x)$ is a strict super-solution of the following problem

$$\begin{cases} w_t - \Delta w = w^p + f(x), & (x, t) \in \Omega \times (0, T), \\ \frac{\partial w}{\partial \nu} + \beta w = 0, & (x, t) \in \partial\Omega \times [0, T], \\ w(x, 0) = h_\beta(x), & x \in \Omega. \end{cases} \quad (6.12)$$

Let $w(x, t)$ be the solution of problem (6.12). Then, $w(x, t)$ exists globally and is strictly decreasing with respect to t . Since $h_\beta(x) > U_\beta(x)$, it follows from the comparison principle that $w(x, t) > U_\beta(x)$ for any $(x, t) \in \Omega \times [0, +\infty)$. Hence, we may assume that

$$\lim_{t \rightarrow +\infty} w(x, t) = W(x).$$

Moreover, it is easy to prove that $W(x)$ satisfies

$$\begin{cases} -\Delta W = W^p + f(x), & x \in \Omega, \\ W > 0, & x \in \Omega, \\ \frac{\partial W}{\partial \nu} + \beta W = 0, & x \in \partial\Omega. \end{cases} \quad (6.13)$$

This will imply $W(x) \equiv U_\beta(x)$. Otherwise, on one hand, we should have $W(x) > U_\beta(x)$, on the other hand, strong comparison principle should imply $W(x) < u_\beta(x)$ since $h_\beta(x) < u_\beta(x)$.

This contradicts lemma 6.4 since $W(x)$ and $u_\beta(x)$ are two solutions of problem (1.1) which are distinct to the minimal solution $U_\beta(x)$ and do not intersect each other. Thus, we have

$$\lim_{t \rightarrow +\infty} w(x, t) = U_\beta(x).$$

Let $v(x, t)$ be the solution of the problem

$$\begin{cases} v_t - \Delta v = v^p + f(x), & (x, t) \in \Omega \times (0, T), \\ \frac{\partial v}{\partial \nu} + \beta v = 0, & (x, t) \in \partial\Omega \times [0, T], \\ w(x, 0) = 0, & x \in \Omega. \end{cases} \quad (6.14)$$

Then, $v(x, t)$ exists globally and is strictly increasing with respect to t . Moreover, comparison principle implies $v(x, t) \leq U_\beta(x)$ since $U_\beta(x)$ is a super-solution of problem (6.14). Consequently, $v(x, t)$ converges when $t \rightarrow +\infty$. It is obvious that

$$\lim_{t \rightarrow +\infty} v(x, t) = U_\beta(x).$$

Since $0 \leq u_0(x) \leq \eta u_\beta(x) < h_\beta(x)$, comparison principle assures that

$$v(x, t) \leq u(x, t; u_0) \leq w(x, t).$$

By applying squeeze principle, we obtain

$$\lim_{t \rightarrow +\infty} u(x, t; u_0) = U_\beta(x).$$

This completes the proof of theorem 1.4 (ii).

Finally, we prove theorem 1.4 (iii) by contradiction. To this end, we assume that $u_0(x) \geq u_\beta(x)$, $u_0(x) \not\equiv u_\beta(x)$, and problem (1.9) has a global solution $u(x, t, u_0)$. By the strong comparison principle, we have

$$u(x, t; u_0) > u_\beta(x) \quad \text{for any } (x, t) \in \overline{\Omega} \times (0, +\infty). \quad (6.15)$$

Hence, we may assume, by replacing $u_0(x)$ with $u(x, T; u_0)$ for some $T > 0$ if necessary, that $u_0(x) \geq \eta u_\beta(x)$ for some $\eta > 1$. Let $H_\beta(x) = \eta(u_\beta(x) - U_\beta(x)) + U_\beta(x)$. Then, by making use of lemma 6.3 with $\eta > 1$, we can verify that $H_\beta(x)$ satisfies

$$\begin{cases} -\Delta H_\beta < H_\beta^p + f(x), & x \in \Omega, \\ H_\beta > 0, & x \in \Omega, \\ \frac{\partial H_\beta}{\partial \nu} + \beta H_\beta = 0, & x \in \partial\Omega. \end{cases} \quad (6.16)$$

This implies that $H_\beta(x)$ is a strict sub-solution of the following problem

$$\begin{cases} G_t - \Delta G = G^p + f(x), & (x, t) \in \Omega \times (0, T), \\ \frac{\partial G}{\partial \nu} + \beta G = 0, & (x, t) \in \partial\Omega \times [0, T], \\ G(x, 0) = H_\beta(x), & x \in \Omega. \end{cases} \quad (6.17)$$

Let $G(x, t)$ be the solution of problem (6.17). Then, it follows from the comparison principle that $G(x, t) \leq u(x, t; u_0)$ for any (x, t) due to $H_\beta(x) < \eta u_\beta(x) \leq u_0(x)$. Consequently, $G(x, t)$ exists globally and is strictly increasing in t . By theorem 4.1, there exists a positive constant M independent of t such that $G(x, t) \leq M$ for any (x, t) . Hence, $G(x, t)$ converges to some function $g(x)$. Obviously, $g(x)$ is a solution of problem (1.1). Noting that $H_\beta(x) > u_\beta(x)$, it follows from the strong comparison principle that $G(x, t) > u_\beta(x)$ for any (x, t) . Consequently, $g(x) > u_\beta(x)$. This contradicts lemma 6.4 since $g(x)$ and $u_\beta(x)$ are two solutions of problem (1.1) which are distinct to the minimal solution $U_\beta(x)$ and do not intersect each other. $\square$

References

- [1] A.Ambrosetti, P.H.Rabinowitz, Dual variation methods in critical point theory and applications, *J.Funct.Anal.*14 (1973) 349-381.
- [2] T.Cazenave, F.Dickstein and M.Escobedo, A semilinear heat equation with concave-convex nonlinearity, *Rendiconti di Matematica Serie VII* 19 (1999) 211-242.
- [3] C.V.Coffman, On the positive solutions of boundary-value problem for a class of nonlinear differential equation, *J.Differential Equations* 3 (1967) 92-111.
- [4] Qiuyi Dai and Yonggeng Gu, Positive solutions for non-homogeneous semilinear elliptic equations with data that changes sign, *Proc.Roy.Edinburgh* 133A (2003) 297-306.
- [5] Qiuyi Dai and Yuxia Fu, Nondegeneracy and uniqueness of positive solutions for Robin problem of second order ordinary differential equations and its applications, *J. Differential Equations* 241 (2007) 32-61.
- [6] Qiuyi Dai and Jianfu Yang, Positive solutions of inhomogeneous elliptic equations with indefinite data, *Nonlinear Analysis* 58 (2004) 571-589.
- [7] L.Damascelli, M.Grossi and F.Pacella, Qualitative properties of positive solutions of semilinear elliptic equations in symmetric domains via the maximum principle, *Ann.Inst.H.poincare Anal.Non Lineaire* 16 (5) (1999) 631-652.
- [8] Yinbin Deng, Yi Li and Fen Yang, On the stability of the positive steady states for a nonhomogeneous semilinear Cauchy problem, *J.Differential Equations* 228 (2006) 507-529.
- [9] Yinbin Deng, Existence of Multiple Positive Solutions for $-\Delta u = \lambda u + u(N+2)/(N-2) + \mu f(x)$, *Acta Math. Sinica* 9(1993) 311-320.
- [10] Yuxia Fu and Qiuyi Dai, Positive solutions of the Robin problem for semilinear elliptic equations on annuli, *Rend.Lincei Mat.Appl* 19 (2008) 175-188.
- [11] B.Gidas and J.Spruck, A priori bounds for positive solution of nonlinear elliptic equations, *Comm. In Partial Differential equations* 6 (1981) 883-901.
- [12] Y.Giga, A bound for global solutions of semilinear heat equations, *Comm. Math. Phys.*, 103 (1986) 415-421.
- [13] Changfeng Gui, Weiming Ni and Xuefeng Wang, On the stability and instability of positive steady states of a semilinear heat equation in R^n , *Comm.Pure Appl.Math* 45(9) (1992) 1153-1181.
- [14] Changfeng Gui, Weiming Ni and Xuefeng Wang, Further Study on a Nonlinear Heat Equation, *J. Differential Equations* 169 (2001) 588-613.
- [15] Yanyan Li, Existence of many positive solutions of semilinear elliptic equations on annulus, *J. Differential Equations* 83 (1990) 348-367.
- [16] S.S.Lin, Existence of positive nonradial solutions for nonlinear elliptic equations in annulus domains, *Trans. Amer. Math. Soc.*, 332(1992), 775-791

- [17] S.S.Lin, Existence of many positive nonradial solutions for nonlinear elliptic equations on annulus, *J. Differential Equations*, 103(1993), 338-349
- [18] Yonggeng Gu and Tong Liu, A prior estimate and existence of positive solutions of semilinear elliptic equation with the third boundary value problem , *Journal of System Science and Complexity*,14(2001) 388-398.
- [19] Li Ma and Ning Su, Existence, multiplicity and stability results for positive solutions of nonlinear p -Laplacian equations, *Chin. Ann. Math.* 25B(2) (2004) 275-286.
- [20] P.Polacik, P.Quittner and P.Souplet, Singularity and decay estimates in superlinear problems via Liouville-type theorems. Part II: Parabolic equations, *Indiana Univ. Math. J.* 56 (2007), 879-908
- [21] P.Polacik, P.Quittner and P.Souplet, Singularity and decay estimates in superlinear problems via Liouville-type theorems. Part I: Elliptic equations and systems. *Duke Math. J.* 139 (2007), 555-579
- [22] G.Tarantello, On nonhomogeneous elliptic equations involving critical Sobolev exponent, *Ann.Inst.Henri Poincaré* 9 (1992) 281-304.
- [23] Xuefeng Wang, On the cauchy problem for reaction-diffusion equations, *Transactions of the American Mathematical society* 337 (1993) 549-590.
- [24] Xujia Wang, Neumann problems of semilinear elliptic equations involving critical Sobolev exponents, *J. Differential Equations*, 93(1991), 283-310
- [25] F.B.Weissler, Existence and non-existence of global solutions for a semilinear heat equation, *Israel Journal of Mathematics* 38(1981) 29-40.